

www.tajpharma.com

Dosage Forms & Strengths

Tablet : 2mg & 5mg

Chronic Lymphatic (Lymphocytic) Leukemia

0.1 mg/kg/day for 3-6 weeks or 0.4 mg/kg

(increased by 0.1 mg/kg/dose until response/toxicity observed) biweekly or

0.4 mg/kg (increased by 0.1 mg/kg/dose until response observed) monthly or

0.03-0.1 mg/kg/day continuously

**Briviran**<sup>®</sup>  
(chlorambucil) 2/5 mg  
Tablets

STORE AT : 2°C TO 8°C. (REFRIGERATE. DO NOT FREEZE). PROTECT FROM LIGHT.  
Provided that the outer coating of the tablet is intact, there is no risk in handling  
Chlorambucil Tablets.  
Chlorambucil Tablets should not be divided.

**Briviran**<sup>®</sup>  
(chlorambucil) 2/5 mg  
Tablets

Briviran<sup>®</sup> Chlorambucil 2mg & 5mg Tablets medication **is used to treat certain types of cancer (such as leukemia, lymphoma)**. Chlorambucil belongs to a class of drugs known as alkylating agents. It works by slowing or stopping the growth of cancer cells.

This medication is used to treat certain types of cancer (such as leukemia, lymphoma). Chlorambucil belongs to a class of drugs known as alkylating agents.

**It works by slowing or stopping the growth of cancer cells.**

**How to use Chlorambucil Tablet**

Take this medication by mouth as directed by your doctor, usually once daily. Unless otherwise directed by your doctor, drink plenty of fluids to help prevent side effects. The dosage is based on your medical condition, weight, and response to treatment. Do not increase your dose or take this medication more often than directed. Your condition will not improve any faster and your risk of side effects will increase.

Since this drug can be absorbed through the skin and lungs and may harm an unborn baby, women who are pregnant or who may become pregnant should not handle this medication or breathe the dust from the tablets.

**CHLORAMBUCIL IS A CANCER MEDICINE THAT INTERFERES WITH THE GROWTH AND SPREAD OF CANCER CELLS IN THE BODY.**

Chlorambucil is used to treat several types of cancer, including Hodgkin's disease and certain types of leukemia or lymphoma.

BRIVIRAN is indicated in the treatment of:-

- Hodgkin's disease
- certain forms of non-Hodgkin's lymphoma
- chronic lymphocytic leukaemia
- Waldenstrom's macroglobulinaemia
- Advanced ovarian adenocarcinoma.

BRIVIRAN has a significant therapeutic effect on a proportion of patients with breast cancer.

**Food and Drug Administration (FDA) approved .Prescription Only (POM)**

A Taj Pharma India Product

**TAJ**<sup>®</sup>  
PHARMA  
A TAJ Enterprise