www.tajpharma.com

The active substance in Cadrol ® Bisoprolol Fumarate Tablets is bisoprolol. Cadrol ® belongs to a group of medicines called beta-blockers. These medicines work by affecting the body's response to some nerve impulses, especially in the heart. As a result, bisoprolol slows down the heart rate and makes the heart more efficient at pumping blood around the body.





Cadrol ® (bisoprolol fumarate) is a synthetic, beta1-selective (cardioselective) adrenoceptor blocking agent.

Cadrol is a drug belonging to the group of beta blockers, a class of medicines used primarily in cardiovascular diseases.

More specifically, it is a selective type \$1 adrenergic receptor blocker.

Cadrol ® (Bisoprolol Fumarate) is used alone or in combination with other medications to treat high blood pressure. Cadrol is in a class of medications called beta blockers. It works by relaxing blood vessels and slowing heart rate to improve and decrease blood pressure. How should this medicine be used?

Cadrol ® (Bisoprolol Fumarate) comes as a tablet to take by mouth. It is usually taken once a day. To help you remember to take Cadrol ® (Bisoprolol Fumarate), take it around the same time every day.

Follow the directions on your prescription label carefully, and ask your doctor or pharmacist to explain any part you do not understand.

Take Cadrol ® (Bisoprolol Fumarate) exactly as directed. Do not take more or less of it or take it more often than prescribed by your doctor.

Your doctor will probably start you on a low dose of Cadrol (Bisoprolol Fumarate) and gradually increase your dose.

Cadrol (Bisoprolol Fumarate) controls high blood pressure but does not cure it. It may take a few weeks before you feel the full benefit of Cadrol. Continue to take Cadrol (Bisoprolol Fumarate) even if

you feel well. Do not stop taking bisoprolol without talking to your doctor. Suddenly stopping bisoprolol may cause angina (chest pain), heart attack, or irregular heartbeat. Your doctor will probably decrease your dose gradually.

The active substance is bisoprolol fumarate.

Each tablet contains of bisoprolol fumarate / The other ingredients are lactose monohydrate, cellulose microcrystalline [E460], magnesium stearate [E572] and crospovidone [E1202].

Bisoprolol fumarate is supplied as 1.5 mg , 5 mg and 10 mg tablets.

DOSAGE AND ADMINISTRATION

The dose of bisoprolol fumarate must be individualized to the needs of the patient. The usual starting dose is 5 mg once daily. In some patients, 2.5 mg may be an appropriate starting dose (see Bronchospastic Disease in WARNINGS). If the antihypertensive effect of 5 mg is inadequate, the dose may be increased to 10 mg and then, if necessary, to 20 mg once daily.