

Candesartan Cilexetil Tablets 16mg/32mg Taj Pharma

Candesartan Cilexetil Tablets

Patient Information leaflet: Important for the user

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- The full name of this medicine is Candesartan 4mg, 8mg, 16mg and 32mg Tablets but within the leaflet it will be referred to as Candesartan tablets.

What is in this leaflet:

- 1 What Candesartan tablets are and what they are used for
- 2 What you need to know before you take
- 3 How to take
- 4 Possible side effects
- 5 How to store
- 6 Contents of the pack and other information

1. What Candesartan tablets are and what they are used for

The active ingredient is candesartan cilexetil. This belongs to a group of medicines called angiotensin II receptor antagonists. It works by making your blood vessels relax and widen. This helps to lower your blood pressure. It also makes

it easier for your heart to pump blood to all parts of your body.

This medicine is used for:

- treating high blood pressure (hypertension) in adult patients and in children and adolescents aged 6<18 years.
- treating adult heart failure patients with reduced heart muscle function when Angiotensin Converting Enzyme (ACE) inhibitors cannot be used or in addition to ACE-inhibitors when symptoms persist despite treatment and mineralocorticoid receptor antagonists (MRA) cannot be used. (ACE-inhibitors and MRAs are medicines used to treat heart failure).

2. What you need to know before you take

Do not take Candesartan tablets if:

- you are **allergic** to candesartan cilexetil or any of the other ingredients of this medicine (listed in section 6).
- you are more than **3 months pregnant** (it is also better to avoid Candesartan tablets in early pregnancy – see pregnancy section).
- you have **severe liver** disease or **biliary obstruction** (a problem with the drainage of the bile from the gall bladder).
- the patient is a child under 1 year of age.
- you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren.

If you are not sure if any of these apply to you, talk to your doctor or pharmacist before taking Candesartan tablets.

Warnings and precautions

Talk to your doctor before taking Candesartan tablets if you:

- have **heart, liver** or **kidney** problems, or are on **dialysis**.
- have recently had a **kidney transplant**.
- are **vomiting**, have recently had severe vomiting, or have **diarrhoea**.
- have a disease of the adrenal gland called **Conn's syndrome** (also called **primary hyperaldosteronism**).
- have **low blood pressure**.
- are taking any of the following medicines used to treat high blood pressure:
 - an ACE-inhibitor (for example enalapril, lisinopril, ramipril) in particular if you have diabetes related kidney problems.
 - aliskiren
- have ever had a **stroke**.
- are taking an ACE-inhibitor together with a medicine which belongs to the class of medicines known as mineralocorticoid receptors antagonists (MRA). These medicines are for the treatment of heart failure (see "Other medicines and Candesartan tablets").

You must tell your doctor if you think that you are (or might become) pregnant. Candesartan tablets is not recommended in early pregnancy and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

Your doctor may check your kidney function, blood pressure and the amount of electrolytes (e.g. potassium) in your blood at regular intervals.

See also information under the heading "Do not take Candesartan tablets if"

If you are going to have an operation, tell your doctor or dentist that you are taking Candesartan tablets. This is because Candesartan tablets, when combined with some anaesthetics, may cause a drop in blood pressure.

Children and adolescents

Candesartan has been studied in children. For more information, talk to your doctor. Candesartan tablets must not be given to children under 1 year of age due to the potential risk to the developing kidneys.

Other medicines and Candesartan tablets

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines. Candesartan tablets can affect the way some other medicines work and some medicines can have an effect on Candesartan tablets. If you are using certain medicines, your doctor may need to do blood tests from time to time.

In particular, tell your doctor if you are using any of the following medicines:

- Other medicines to help lower your blood pressure, including beta-blockers, diazoxide and ACE inhibitors such as enalapril, captopril, lisinopril or ramipril.
- Non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen, naproxen, diclofenac, celecoxib or etoricoxib (medicines to relieve pain and inflammation).
- Aspirin (if you are taking more than 3g each day) (medicine to relieve pain and inflammation).
- Potassium supplements or salt substitutes containing potassium (medicines that increase the amount of potassium in your blood).
- Heparin (a medicine for thinning the blood).
- Water tablets (diuretics).
- Lithium (a medicine for mental health problems).

Your doctor may need to change your dose and/or to take other precautions:

- If you are taking an ACE-inhibitor or aliskiren (see also information under the headings "Do not take Candesartan tablets if" and "Warnings and precautions").

- If you are being treated with an ACE-inhibitor together with certain other medicines to treat your heart failure, which are known as mineralocorticoid receptors antagonists (MRA) (for example spironolactone, eplerenone).

Candesartan tablets with food and drink

- You can take Candesartan tablets with or without food.
- When you are prescribed Candesartan tablets, discuss with your doctor before drinking alcohol. Alcohol may make you feel faint or dizzy.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Candesartan tablets before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Candesartan tablets. Candesartan tablets is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

Breast-feeding

Tell your doctor if you are breast-feeding or about to start breast-feeding. Candesartan tablets are not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn, or was born prematurely.

Driving and using machines

Some people may feel tired or dizzy when taking Candesartan tablets. If this happens to you, do not drive or use any tools or machines.

Candesartan tablets contain lactose

Candesartan tablets contain lactose which is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. It is important to keep taking Candesartan tablets every day.

You can take Candesartan tablets with or without food.

Swallow the tablet with a drink of water.

Try to take the tablet at the same time each day. This will help you to remember to take it.

High blood pressure:

- The recommended dose of Candesartan tablets is 8mg once a day. Your doctor may increase this dose to 16mg once a day and further up to 32mg once a day depending on blood pressure response.
- In some patients, such as those with liver problems, kidney problems or those who recently have lost body fluids, e.g., through vomiting or diarrhoea or by using water tablets, the doctor may prescribe a lower starting dose.
- Some black patients may have a reduced response to this type of medicine, when given as the only treatment, and these patients may need a higher dose.

Heart failure:

- The recommended starting dose of Candesartan tablets is 4mg once a day. Your doctor may increase your dose by doubling the dose at intervals of at least 2 weeks up to

32mg once a day. Candesartan tablets can be taken together with other medicines for heart failure, and your doctor will decide which treatment is suitable for you.

Use in children and adolescents with high blood pressure

Children 6 to <18 years of age: The recommended starting dose is 4mg once a day.

For patients weighing <50kg: In some patients whose blood pressure is not adequately controlled, your doctor may decide the dose needs to be increased to a maximum of 8mg once a day.

For patients weighing ≥50kg: In some patients whose blood pressure is not adequately controlled, your doctor may decide the dose needs to be increased to 8mg once a day or to 16mg once a day.

If you take more Candesartan tablets than you should

If you take more Candesartan tablets than prescribed by your doctor, contact a doctor or pharmacist immediately for advice.

If you forget to take Candesartan tablets

Do not take a double dose to make up for a forgotten tablet. Just take the next dose as normal.

If you stop taking Candesartan tablets

If you stop taking Candesartan tablets, your blood pressure may increase again. Therefore do not stop taking Candesartan tablets without first talking to your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. It is important that you are aware of what these side effects may be.

Stop taking Candesartan tablets and seek medical help immediately if you have any of the following allergic reactions:

- difficulties in breathing, with or without swelling of the face, lips, tongue and/or throat.
- swelling of the face, lips, tongue and/or throat, which may cause difficulties in swallowing.
- severe itching of the skin (with raised lumps).

Candesartan tablets may cause a reduction in number of white blood cells. Your resistance to infection may be decreased and you may notice tiredness, an infection or a fever. If this happens contact your doctor.

Your doctor may occasionally do blood tests to check whether Candesartan tablets has had any effect on your blood (agranulocytosis).

Other possible side effects include:

Common (may affect up to 1 in 10 users)

- Feeling dizzy/spinning sensation.
- Headache.
- Respiratory infection.
- Low blood pressure. This may make you feel faint or dizzy.
- Changes in blood test results:
 - An increased amount of potassium in your blood, especially if you already have kidney problems or heart failure. If this is severe you may notice tiredness, weakness, irregular heart beat or pins and needles.
- Effects on how your kidneys work, especially if you already have kidney problems or heart failure. In very rare cases, kidney failure may occur.

Very rare (may affect up to 1 in 10,000 users)

- Swelling of the face, lips, tongue and/or throat.

- A reduction in your red or white blood cells. You may notice tiredness, an infection or a fever.
- Skin rash, lumpy rash (hives).
- Itching.
- Back pain, pain in joints and muscles.
- Changes in how your liver is working, including inflammation of the liver (hepatitis). You may notice tiredness, yellowing of your skin and the whites of your eyes and flu like symptoms.
- Nausea.
- Changes in blood test results:
- A reduced amount of sodium in your blood. If this is severe then you may notice weakness, lack of energy, or muscle cramps.
- Cough.

In children treated for high blood pressure, side effects appear to be similar to those seen in adults, but they happen more often. Sore throat is a very common side effect in children but not reported in adults and runny nose, fever and increased heart rate are common in children but not reported in adults.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

5. How to store

Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the carton and on the blister pack after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines no longer used. These measures will help protect the environment.

6. Contents of the pack and other information

What Candesartan tablets contains

a) Each Uncoated tablet contains:
Candesartan Cilexetil Tablets USP 16mg
Excipients: q.s

b) Each Uncoated tablet contains:
Candesartan Cilexetil Tablets USP 32mg
Excipients: q.s

The other ingredients are lactose monohydrate, maize starch, hydroxypropylcellulose, croscarmellose sodium, magnesium stearate and triethyl citrate.

Available Pack Size: 7, 14, 28, 30, 50, 100 and 500 tablets.

7. Manufactured in India by: TAJ PHARMACEUTICALS LTD.

Mumbai, India

Unit No. 214.Old Bake House,
Maharashtra chambers of Commerce Lane,
Fort, Mumbai - 400001
at:Gujarat, INDIA.

Customer Service and Product Inquiries:
1-800-TRY-FIRST (1-800-222-434 & 1-
800-222-825)

Monday through Saturday 9:00 a.m. to 7:00
p.m. EST

E-mail: tajgroup@tajpharma.com