## www.tajpharma.com

Cifelyra 50 mg daily is indicated for use in combination therapy with a luteinizing hormone-releasing hormone (LHRH) analog for the treatment of Stage D2 metastatic carcinoma of the prostate.

Cifelyra 150 mg daily is not approved for use alone or with other treatments Dosage Forms And Strength

Cifelyra® (bicalutamide) 50 mg tablets for oral administration.

Store at controlled room temperature, 20°-25°C (68°-77°F).





Cifelyra ® Bicalutamide Tablets is used to treat prostate cancer that has spread to other areas of the body. It is used in combination with hormone treatment.

This medication works by blocking the action of male hormones in the prostate, slowing growth of the tumors. This medication should not be used in women or children.

Bicalutamide is a hormonal therapy drug used to treat prostate cancer. It's best to read this with our general information about prostate cancer. You have bicalutamide as tablets. You usually have it as an outpatient. Your cancer doctor or nurse will tell you how often you will have it.

Like all hormonal therapy drugs, bicalutamide can cause side effects. Some of these can be serious, so it's important that you read the detailed information below. How hormonal therapy affects people varies from person to person. Your doctor or nurse can talk to you more about this and give you advice on how to manage any side effects.

Tell your doctor or nurse straight away if you feel unwell or have severe side effects, including any we don't mention here. If you need to see a health professional for any reason other than cancer, always tell them that you are having this treatment.

Qualitative and quantitative composition Each tablet contains 50 mg of bicalutamide.

Excipients with known effect: One tablet contains 60.44 mg lactose monohydrate

Treatment of advanced prostate cancer in combination with Luteinising Hormone-Releasing Hormone (LHRH) analogue therapy or surgical castration.

USES: Bicalutamide is used to treat prostate cancer that has spread to other areas of the body. It is used in combination with hormone treatment. This medication works by blocking the action of male hormones in the prostate, slowing growth of the tumors. This medication should not be used in women or children.

HOW TO USE: Take this medication by mouth, usually once a day in the morning or evening, with or without food, or as directed by your doctor. Use this medication regularly in order to get the most benefit from it. Remember to use it at the same time each day. The dosage is based on your medical condition and response to therapy. Do not increase your dose or take this medication more often without your doctor's approval. Your condition will not improve any faster, and the risk of serious side effects may be increased. Since this drug can be absorbed through the skin, women who are pregnant or who may become pregnant should not handle this medication.

Food and Drug Administration (FDA) approved .Prescription Only (POM)

PHARMA
A TAJ Enterprise