

## **Exemestane Tablets USP 25mg Taj Pharma**

### **Package leaflet: Information for the user**

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it
- If you have any further questions, ask your doctor, pharmacist or
- This medicine has been prescribed for you Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section

### **What is in this leaflet:**

1. What Exemestane Taj Pharma tablet is and what it is used for
2. What you need to know before you take Exemestane Taj Pharma tablets
3. How to take Exemestane Taj Pharma tablets
4. Possible side effects
5. How to store Exemestane Taj Pharma tablets
6. Contents of the pack and other information

### **1. WHAT EXEMESTANE TAJ PHARMA TABLET IS AND WHAT IT IS USED FOR**

Exemestane Taj Pharma belongs to a group of medicines known as aromatase inhibitors. These medicines interfere with a substance called aromatase, which is needed to make the female sex hormones, oestrogens,

especially in postmenopausal women. Reduction in oestrogen levels in the body is a way of treating hormone dependent breast cancer.

Exemestane Taj Pharma is used to treat hormone dependent early breast cancer in postmenopausal women after they have completed 2-3 years of treatment with the medicine tamoxifen.

Exemestane Taj Pharma is also used to treat hormone dependent advanced breast cancer in postmenopausal women when a different hormonal drug treatment has not worked well enough.

### **2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE EXEMESTANE TAJ PHARMA TABLETS**

#### **Do not take Exemestane Taj Pharma tablets**

- if you are or have previously been allergic to Exemestane Taj Pharma or any of the other ingredients of this medicine (listed in section 6).
- if you have not already been through ‘the menopause’, e. you are still having your monthly period,
- if you are pregnant, likely to be pregnant or

### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Exemestane Taj Pharma tablets.

- Before treatment with Exemestane Taj Pharma, your doctor may want to take blood samples to make sure you have reached the

- Routine checking of your vitamin D level will also be made before treatment, as your level may be very low in the early stages of breast You will be given vitamin D supplement if your levels are below normal.
- Before taking Exemestane Taj Pharma, tell your doctor if you have problems with your liver or
- Tell your doctor if you have a history or are suffering from any condition which affects the strength of your Your doctor may want to measure your bone density before and during the treatment of Exemestane Taj Pharma. This is because medicines of this class lower the levels of female hormones and this may lead to a loss of the mineral content of bones, which might decrease their strength.

#### **Other medicines and Exemestane Taj Pharma Tablets**

Please tell your doctor if you are taking or have recently taken or might take any other medicines, including medicines obtained without a prescription.

Exemestane Taj Pharma tablets should not be given at the same time as hormone replacement therapy (HRT).

The following medicines should be used cautiously when taking Exemestane Taj Pharma tablets. Let your doctor know if you are taking medicines such as:

- rifampicin (an antibiotic),
- carbamazepine or phenytoin (anticonvulsants used to treat epilepsy),
- the herbal remedy St John's Wort, (Hypericum perforatum), or preparations containing

#### **Exemestane Taj Pharma Tablets with food, drink and alcohol**

Exemestane Taj Pharma Tablets should be taken after a meal at approximately the same time each day.

#### **Pregnancy, breast-feeding and fertility**

Do not take Exemestane Taj Pharma tablets if you are pregnant or breastfeeding. If you are pregnant or think you might be, tell your doctor.

Discuss contraception with your doctor if there is any possibility that you may become pregnant. Ask your doctor or pharmacist for advice before taking any medicine.

#### **Driving and using machines**

If you feel drowsy, dizzy or weak whilst taking Exemestane Taj Pharma tablets, you should not attempt to drive or operate machinery.

### **3. HOW TO TAKE EXEMESTANE TAJ PHARMA TABLETS**

#### **Adults and the elderly**

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

Exemestane Taj Pharma tablets should be taken by mouth after meal at approximately the same time each day. Your doctor will tell you how to take Exemestane Taj Pharma tablets and for how long.

The recommended dose is one 25mg tablet daily.

If you need to go to the hospital whilst taking Exemestane Taj Pharma tablets, let the medical staff know what medication you are taking.

#### **Use in children**

Exemestane Taj Pharma tablets is not suitable for use in children.

### **If you take more Exemestane Taj Pharma Tablets than you should**

If too many tablets are taken by accident, contact your doctor at once or go straight to the nearest hospital casualty department. Show them the pack of Exemestane Taj Pharma tablets.

### **If you forget to take Exemestane Taj Pharma Tablets**

Do not take a double dose to make up for a forgotten tablet.

If you forget to take your tablet, take it as soon as you remember. If it is nearly time for the next dose, take it at the usual time.

### **If you stop taking Exemestane Taj Pharma Tablets**

Do not stop taking your tablets even if you are feeling well, unless your doctor tells you.

If you have any further questions on the use of medicine, ask your doctor, pharmacist or nurse.

## **4. POSSIBLE SIDE EFFECTS**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Exemestane Taj Pharma tablets is well tolerated and the following side effects observed in patients treated with Exemestane Taj Pharma tablets are mainly mild or moderate in nature. Most of the side effects are associated with a shortage of oestrogen (e.g. hot flushes).”

Hypersensitivity, inflammation of the liver (hepatitis) and inflammation of the bile

ducts of the liver which cause yellowing of the skin (cholestatic hepatitis) may occur. Symptoms include feeling generally unwell, nausea, jaundice (yellowing of the skin and eyes), itching, right sided abdominal pain and loss of appetite. Contact your doctor promptly to seek urgent medical advice if you think you have any of these symptoms.

### **Very common side effects (may affect more than 1 in 10 people):**

- Depression
- Difficulty sleeping
- Headache
- Hot flushes
- Dizziness
- Feeling sick
- Increased sweating
- Muscle and joint pain (including osteoarthritis, back pain, arthritis and joint stiffness)
- Tiredness
- A reduction in the number of white blood cells
- Abdominal pain
- Elevated level of liver enzymes
- Elevated level of a haemoglobin breakdown in the blood
- Elevated level of a blood enzyme in the blood due to liver damage
- Pain

### **Common side effects (may affect up to 1 in 10 people):**

- Loss of appetite Carpal tunnel syndrome (a combinations of pins and needles, numbness and pain affecting all of the hand except the little finger) or tingling/prickling of the skin
- Stomach ache ,vomiting (being sick), constipation, indigestion, diarrhoea
- Hair loss
- Skin rash, hives and itchiness

- Thinning of bones which might decrease their strength (osteoporosis), leading to bone fractures (breaks or cracks) in some cases
- Pain, swollen hands and feet
- A reduction in the number of platelets in the blood
- Muscle weakness

**Uncommon side effects (may affect up to 1 in 100 people):**

- Hypersensitivity

**Rare side effects (may affect up to 1 in 1,000 people):**

- A breakout of small blisters on an area of the skin in a rash
- Drowsiness
- Inflammation of the liver
- Inflammation of the bile ducts of the liver which cause yellowing of the skin

Not Known (frequency cannot be estimated from the available data)

- Low level of certain white blood cells in the blood

Changes in the amount of certain blood cells (lymphocytes) and platelets circulating in your blood, especially in patients with a pre-existing lymphopenia (reduced lymphocytes in the blood) may also be seen.

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

**5. HOW TO STORE EXEMESTANE TAJ PHARMA TABLETS**

- Keep this medicine out of the sight and reach of

- Do not use this medicine after the expiry date which is stated on the outer carton and blister after EXP. The expiry date refers to the last day of that
- This medicine does not require any special storage
- Do not throw away any medicines via wastewater or household Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

**6. CONTENTS OF THE PACK AND OTHER INFORMATION**

**What Exemestane Taj Pharma Tablets contains**

The active substance is Exemestane Taj Pharma. Each film-coated tablet contains Exemestane Taj Pharma 25mg

The other ingredients are:

Mannitol, cellulose microcrystalline, crospovidone, sodium starch glycolate, hypromellose, polysorbate 80, colloidal anhydrous silica, magnesium stearate, Hypromellose 6cp titanium dioxide macrogol

**What Exemestane Taj Pharma tablets looks like and contents of the pack**

Exemestane Taj Pharma 25mg Tablets are available in the White opaque PVC/PVdC-Alu blister packs of 15, 20, 28, 30, 60, 90, 98, 100 and 120 tablets.

Not all pack sizes may be marketed.

**Manufactured in India By:**

**TAJ PHARMACEUTICALS LIMITED**  
at SURVEY NO.188/1 TO 189/1,190/1 TO 4, ATHIYAWAD, DABHEL, DAMAN-396210 (INDIA)