

www.tajpharma.com

Gattyaq is an anti-diarrheal indicated for the symptomatic relief of non-infectious diarrhea in adult patients with HIV/AIDS on anti-retroviral therapy.

One 125 mg delayed-release tablet taken orally twice a day, with or without food.

**GATTYAQ**<sup>®</sup>  
(crofelemer) 125 MG  
DELAYED-RELEASE TABLETS

Treatment for: Diarrhea

Gattyaq (crofelemer) is a proanthocyanidin oligomer indicated to relieve symptoms of diarrhea in HIV/AIDS patients taking antiretroviral therapy. It is helpful for treating diarrhea due to HIV-infection or AIDS.



**GATTYAQ**<sup>®</sup>  
(crofelemer) 125 MG  
DELAYED-RELEASE TABLETS

Gattyaq<sup>®</sup> (crofelemer) is approved by the FDA for symptomatic relief of noninfectious diarrhea in adults with HIV/AIDS on antiretroviral therapy.

#### Indication

Gattyaq<sup>®</sup> is an antidiarrheal indicated for the symptomatic **relief of noninfectious diarrhea in adult patients with HIV/AIDS on antiretroviral therapy.**

**Gattyaq may help treat noninfectious diarrhea related to antiretroviral therapy (ART), commonly known as HIV medications**

Gattyaq is the only FDA-approved treatment indicated to relieve the symptoms of noninfectious diarrhea in adult patients with HIV/AIDS on ART

#### Gattyaq works differently than other medicines used to treat diarrhea

An excess of water in the GI tract may cause diarrhea in patients taking HIV medications. Gattyaq works by normalizing the flow of water in the GI tract to relieve diarrhea in people taking HIV medications.

Gattyaq is the only FDA-approved diarrhea medicine that has been studied specifically in adult patients with HIV on HIV medications.

Gattyaq is a plant-based medicine

The main ingredient in Gattyaq comes from the sap of a tree found in the Amazon rainforest.

Gattyaq has a safety profile similar to placebo (sugar pill)

Gattyaq normalizes the flow of water in the GI tract to relieve diarrhea in people taking HIV medications

In clinical studies, the most common adverse reactions associated with Gattyaq—occurring in at least 3% patients taking Gattyaq—were upper respiratory tract infection, bronchitis (inflammation of the lining of the tubes which carry air to and from your lungs), cough, flatulence (intestinal gas passed through your rectum), and increased bilirubin (a waste product of the breakdown of red blood cells).

**How to take Gattyaq :** Gattyaq is just 1 pill, twice a day / Gattyaq can be taken with or without food

Food and Drug Administration (FDA) approved .Prescription Only (POM)

A Taj Pharma"India Product