

www.tajpharma.com

MacriStar® Triptorelin **better than surgical castration**
in lowering testosterone levels

MacriStar
TRIPTORELIN FOR INJECTION **3.75 MG**

**The MacriStar® brand of triptorelin is used in men to
treat the symptoms of prostate cancer.**

Store MacriStar vial supplied with MIXJECT Dose Delivery System (with Sterile Water for Injection) at 20-25°C; excursions permitted: 15 – 30 °C.
Store MacriStar vial (without Sterile Water for Injection) at 20-25°C; excursions permitted: 15 - 30°C.
Do not freeze. Protect from light.
Keep out of reach of children.

MacriStar
TRIPTORELIN FOR INJECTION **3.75 MG**

MacriStar (triptorelin for injectable suspension) is indicated for:
- The management and relief of chronic pain associated with endometriosis.

MacriStar belongs to a class of drugs called gonadotropinreleasing hormone (GnRH) agonists.

What the medication is used for:

Your doctor has prescribed MacriStar to **treat your endometriosis. Endometriosis is a disease in which abnormal tissue grows in the abdomen and other places in the body. It can be treated with pain medication, hormones and surgery.**

What it does:

MacriStar works like a hormone and reduces the level of estrogen in your body. A reduction in the level of your estrogen may help reduce the pain and other symptoms of endometriosis.

The duration of your treatment will be up to 6 months. Once you stop treatment, the level of estrogen will return to normal. Even though you will be stopping treatment after 6 months, you will still benefit from the treatment for many months afterwards.

When it should not be used:

MacriStar should not be used if:

- You are allergic or oversensitive to triptorelin, or to drugs called LHRH agonists, or to any ingredients in the formulation or component of the vial.
- You are or may become pregnant, or if you are nursing. In pregnant women, MacriStar may cause harm to the baby.
- You have abnormal vaginal bleeding that has not been checked by your doctor.
- You are a woman under 18 years of age. MacriStar was not studied in children.

MacriStar must be administered under the supervision of a physician.

Experience in women has been limited to women 18 years of age or older treated for 6 months.

Geriatrics (> 65 years of age): No data are available.

Pediatrics (< 18 years of age): Safety and effectiveness of MacriStar in women with endometriosis under the age of 18 years have not been established.

Usual Dose: Your doctor will administer one injection of MacriStar, containing 3.75 mg triptorelin, into your muscle on a specified day, generally once every 28 days, for a period of 6 months.

Missed Dose: If you forget to have MacriStar administered on the specified day, have it administered as soon as you can.

Food and Drug Administration (FDA) approved .Prescription Only (POM)

A Taj Pharma"India Product

**DO NOT THROW AWAY ANY MEDICINES VIA
WASTEWATER OR HOUSEHOLD WASTE.
ASK YOUR PHARMACIST HOW TO THROW
AWAY MEDICINES YOU NO LONGER USE.
THESE MEASURES WILL HELP PROTECT
THE ENVIRONMENT.**

TAJ
PHARMA®
A TAJ Enterprise