

MAZLERO® can help reduce the risk of stroke in people with atrial fibrillation (AFib) not caused by a heart valve problem and protect against blood clots after knee or hip replacement surgery.

MAZLERO® is a prescription medicine used to reduce the risk of stroke and blood clots in people with atrial fibrillation, not caused by a heart valve problem. For patients currently well managed on warfarin, there is limited information on how MAZLERO® and warfarin compare in reducing the risk of stroke. MAZLERO® is also a prescription medicine used to treat deep vein thrombosis and pulmonary embolism, and to help reduce the risk of these conditions occurring again.

MAZLERO® is also a prescription medicine used to reduce the risk of forming a blood clot in the legs and lungs of people who have just had knee or hip replacement surgery.

IMPORTANT SAFETY INFORMATION

WHAT IS THE MOST IMPORTANT INFORMATION I SHOULD KNOW ABOUT MAZLERO®?

For people taking MAZLERO® for atrial fibrillation:

People with atrial fibrillation (an irregular heart beat) are at an increased risk of forming a blood clot in the heart, which can travel to the brain, causing a stroke, or to other parts of the body. MAZLERO® lowers your chance of having a stroke by helping to prevent clots from forming. If you stop taking MAZLERO®, you may have increased risk of forming a clot in your blood.

Do not stop taking MAZLERO® without talking to the doctor who prescribes it for you. Stopping MAZLERO® increases your risk of having a stroke. If you have to stop taking MAZLERO®, your doctor may prescribe another blood thinner medicine to prevent a blood clot from forming. MAZLERO® can cause bleeding, which can be serious, and rarely may lead to death. This is because MAZLERO® is a blood thinner medicine that reduces blood clotting. While you take MAZLERO® you are likely to bruise more easily and it may take longer for bleeding to stop.