

www.tajpharma.com

NAXILODEX® is a proton pump inhibitor (PPI) indicated for:

- Healing of all grades of erosive esophagitis (EE).
- Maintaining healing of EE and relief of heartburn.
- Treating heartburn associated with symptomatic non-erosive gastroesophageal reflux disease (GERD).

NAXILODEX®
(Dexlansoprazole
Delayed Release Capsules)
30mg & 60mg

NAXILODEX® provides up to 24 hours of relief from heartburn due to GERD.

NAXILODEX®
(Dexlansoprazole
Delayed Release Capsules)
30mg & 60mg

NAXILODEX® (Dexlansoprazole Delayed Release Capsules) is a proton pump inhibitor. Chemically, it is an enantiomer of lansoprazole. NAXILODEX use in the treatment and maintenance of patients with erosive oesophagitis and non-erosive gastro-oesophageal reflux disease (GERD or GORD).

Uses of NAXILODEX® (dexlansoprazole)

Persistent heartburn two or more days a week, despite treatment and diet changes, could be gastroesophageal reflux disease (GERD), also known as acid reflux disease (ARD). Prescription NAXILODEX capsules are used in adults for 4 weeks to treat heartburn related to GERD, for up to 8 weeks to heal acid-related damage to the lining of the esophagus (called erosive esophagitis or EE), and for up to 6 months to continue healing of EE and relief of heartburn. Individual results may vary. Most damage (erosions) heals in 4–8 weeks.

NAXILODEX has a second release of medicine that arrives later in the day

NAXILODEX is a gastroesophageal reflux disease (GERD) treatment that gives you two releases of medicine in one pill. The first release occurs within an hour of taking it, **the second arrives 4–5 hours later.**

How NAXILODEX® works

When you eat, millions of tiny pumps in your stomach produce acid to help digest your food. But sometimes the valve between your stomach and esophagus (called the lower esophageal sphincter or LES) doesn't work quite right, letting stomach acid flow back into your esophagus. This can cause heartburn, a symptom of GERD.

Important Safety Information

Symptom relief does not rule out other serious stomach conditions.

The most common side effects of NAXILODEX® were diarrhea (4.8%), stomach pain (4.0%), nausea (2.9%), common cold (1.9%), vomiting (1.6%), and gas (1.6%).

Dose: Take NAXILODEX any time of day, with or without food

"FOR EROSIVE ESOPHAGITIS, GASTROESOPHAGEAL REFLUX DISEASE"

Food and Drug Administration (FDA) approved .Prescription Only (POM)

A Taj Pharma"India Product