Otipcensa® Carboplatin Injection (carboplatin) belongs to a group of anticancer medicines known as platinum complexes. It is used to treat ovarian cancer.

Otipcensa® Carboplatin is used to treat various types of cancer. It is a chemotherapy drug that contains platinum. It is used alone or in combination with other medications to slow or stop cancer cell growth.



## LIFE WILL WIN AT TAJ PHARMA INDIA.



Carboplatin 10 mg/ml Intravenous Infusion ANTINEOPLASTIC AGENT INDICATED IN THE TREATMENT OF:

- OVARIAN CARCINOMA OF EPITHELIAL ORIGIN
- SMALL CELL LUNG CARCINOMA.
- 5 ml, 15 ml, 45 ml or 60 ml of a sterile solution of carboplatin 10 mg per ml.

Dose (mg) = target AUC (mg/ml x min) x [GFR ml/min + 25]



Carboplatin is used to treat ovarian cancer.

Carboplatin is also used for other types of cancer, including lung, head and neck, endometrial, esophageal, bladder, breast, and cervical; central nervous system or germ cell tumors; osteogenic sarcoma; and as preparation for a stem cell or bone marrow transplant.

Carboplatin (Otipcensa®) 450 mg Injection is an anticancer agent which is used in the treatment of cancer of the female reproductive system. It should be used under the supervision of a qualified physician. Close monitoring of kidney function and blood cell count are necessary while receiving this medication. Carboplatin is indicated in the treatment of:

- ADVANCED STAGE OVARIAN CANCER OF EPITHELIAL ORIGIN
- SMALL CELL LUNG CARCINOMA
- CARCINOMA OF THE HEAD AND NECK
- CARCINOMA OF THE TESTIS
- PAEDIATRIC CEREBRAL TUMOURS
- SOFT TISSUE SARCOMA
- NEUROBLASTOMA

Nature and contents of container Strength Packs 50 mg/5mL 1 x 5 mL glass vial 150 mg/15mL 1 x 20 mL glass vial 450 mg/45mL 1 x 50 mL glass vial 600 mg/60 mL 1 x 100 mL glass vial DO NOT THROW AWAY ANY MEDICINES VIA WASTEWATER OR HOUSEHOLD WASTE. ASK YOUR PHARMACIST HOW TO THROW AWAY MEDICINES YOU NO LONGER USE. THESE MEASURES WILL HELP PROTECT THE ENVIRONMENT.

CAUTION: CARBOPLATIN INJECTION BP IS A POTENT DRUG AND SHOULD BE USED ONLY BY PHYSICIANS EXPERIENCED WITH CANCER CHEMOTHERAPEUTIC DRUGS (SEE WARNINGS AND PRECAUTIONS). BLOOD COUNTS AS WELL AS RENAL AND HEPATIC FUNCTION TESTS MUST BE DONE REGULARLY. DISCONTINUE THE DRUG IF ABNORMAL DEPRESSION OF BONE MARROW OR ABNORMAL RENAL OR HEPATIC FUNCTION IS SEEN.

