

www.tajpharma.com

GASTROPROKINETIC AGENT AND ANTIULCER AGENT

Pintapro® Cinitapride prescribed to treat conditions caused by an overactive digestive tract.

Pintapro®XR

Cinitapride hydrogen tartrate **1 & 3 mg**
EXTENDED RELEASE CAPSULES

Pintapro®XR

Cinitapride hydrogen tartrate **1 & 3 mg**
EXTENDED RELEASE CAPSULES

Pintapro® Cinitapride is a gastroprokinetic agent and antiulcer agent of the benzamide class which is marketed in India, Mexico, Pakistan and Spain. It acts as an agonist of the 5-HT1 and 5-HT4 receptors and as an antagonist of the 5-HT2 receptors.

Pintapro® Cinitapride 1 mg for each dose, 3 mg/daily, for 4 weeks.

Pintapro® Cinitapride is indicated for the treatment of gastrointestinal disorders associated with motility disturbances such as gastroesophageal reflux disease, non-ulcer dyspepsia and delayed gastric emptying.

Pintapro® (Cinitapride) is a antiemetic medication prescribed to treat conditions caused by an overactive digestive tract.

It slows the actions of the muscles to reduce the symptoms of conditions such as acid reflux, ulcer dyspepsia, and delayed gastric emptying.

Uses : Pintapro® (Cinitapride) is a antiemetic medication prescribed to treat conditions caused by an overactive digestive tract.

It slows the actions of the muscles to reduce the symptoms of conditions such as acid reflux, ulcer dyspepsia, and delayed gastric emptying.

Your doctor may also prescribe it to treat other conditions that are not listed here.

Dosage and Administration:

Strictly use this medication as prescribed and follow all instructions provided by your doctor. Safe, suitable, and optimum dosage can vary and is dependent on the patient's health and medical history, as well as the condition you are treating. Pintapro® Cinitapride is typically taken 3 times each day fifteen minutes before each meal. Your treatment will be determined by your health condition, symptoms, and response to previous treatments. Do not alter these directions or self-medicate to prevent unexpected complications.

When taking the doses, swallow the tablets whole along with a full glass of water. Always take the medication on an empty stomach as it should be administered before each of your meals to let it begin to operate before the symptoms occur. Never alter the form of the medication by splitting, chewing, or dissolving it in water as this may destroy it or change its effects.

Food and Drug Administration (FDA) approved .Prescription Only (POM)

A Taj Pharma"India Product