www.tajpharma.com

Primasan® I.V. (Imipenem and Cilastatin for Injection) is a sterile formulation of imipenem (a thienamycin antibiotic) and cilastatin sodium (the inhibitor of the renal dipeptidase, dehydropeptidase I), with sodium bicarbonate added as a buffer. Primasan® I.V. is a potent broad spectrum antibacterial agent for intravenous administration

Primasan ®I.V.

(Imipenem and Cilastatin) for Injection
POWDER FOR SOLUTION FOR INFUSION
[1GM] 500 MG/500 MG

Princisch (Imipenem and Cilastatin) for Injection POWDER FOR SOLUTION FOR INFUSION [1GM] 500 MG/500 MG

Primasan® I.V. (Imipenem and Cilastatin for Injection) is an antibiotic useful for the treatment of a number of bacterial infections.

It is a broad spectrum beta-lactam containing equal quantities of imipenem and cilastatin.

It is related to the penicillin/cephalosporin family of antibiotics but is classified as belonging to the carbapenem class.

Primasan® I.V. (Imipenem and Cilastatin for Injection) is indicated in: lower respiratory tract infections (IV, IM), urinary tract infections (IV), intra-abdominal infections (IV, IM), gynecologic infections (IV, IM), bacterial septicemia (IV), bone and joint infections (IV), skin and skin structure infections (IV, IM), endocarditis (IV) and polymicrobic infections (IV).

Primasan® I.V. (Imipenem and Cilastatin for Injection), an antibiotic, to help treat your infection. The drug will be either injected into a large muscle (such as your buttock or thigh) or added to an intravenous fluid that will drip through a needle or catheter placed in your vein for about 30 minutes, two to four times a day.

The combination of imipenem and cilastatin eliminates bacteria that cause many kinds of infections, including pneumonia and gynecological, skin, stomach, blood, bone, joint, urinary tract, and heart valve infections. This medication is sometimes prescribed for other uses; ask your doctor or pharmacist for more information.

Primasan® I.V.

(IMIPENEM AND CILASTATIN FOR INJECTION)

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Primasan®

I.V. and other antibacterial drugs, Primasan® I.V. should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

Pack size: 1g vial per carton

Primasan ® I.V.

(Imipenem and Cilastatin) for Injection Powder for Solution for Infusion [1GM] 500 MG/500 MG