

www.tajpharma.com

Ryanotar® 50 mg powder for solution for infusion
Each 5 ml Ryanotar vial contains 50 mg of tigecycline. After reconstitution,
1 ml contains 10 mg of tigecycline.
Powder for solution for infusion (powder for infusion).
Orange cake or powder.



RYANOTAR®
(tigecycline) Injection

**LIFE WILL WIN
AT TAJ PHARMA INDIA.**



THE RECOMMENDED DOSAGE REGIMEN FOR RYANOTAR® TIGECYCLINE FOR INJECTION IS AN INITIAL DOSE OF 100 MG, FOLLOWED BY 50 MG EVERY 12 HOURS. INTRAVENOUS INFUSIONS OF TIGECYCLINE FOR INJECTION SHOULD BE ADMINISTERED OVER APPROXIMATELY 30 TO 60 MINUTES EVERY 12 HOURS.



RYANOTAR®
(tigecycline) Injection

Tigecycline is administered intravenously and therefore has 100 % bioavailability.

Tygacil is indicated in adults and in children from the age of eight years for the treatment of the following infections:

- Complicated skin and soft tissue infections (cSSTI), excluding diabetic foot infections ;
 - Complicated intra-abdominal infections (cIAI).
- Tygacil should be used only in situations where other alternative antibiotics are not suitable.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

DO NOT THROW AWAY ANY MEDICINES VIA WASTEWATER OR HOUSEHOLD WASTE. ASK YOUR PHARMACIST HOW TO THROW AWAY MEDICINES YOU NO LONGER USE. THESE MEASURES WILL HELP PROTECT THE ENVIRONMENT.

CONTACT YOUR HEALTH CARE PROVIDER IMMEDIATELY, DAY OR NIGHT, IF YOU SHOULD EXPERIENCE ANY OF THE FOLLOWING SYMPTOMS: FEVER OF 100.4° F (38° C), CHILLS (POSSIBLE SIGNS OF INFECTION) URINARY RETENTION (INABILITY TO URINATE)

Ryanotar® Tigecycline for Injection USP 50mg

Ryanotar 50 MG Injection is an antibiotic which is used to treat infections in the abdomen, skin, and lungs.

This medicine is not used regularly unless necessary. This medicine may cause teeth discoloration in the children. It may increase the risk of sunburns when exposed to the sunlight.

Uses of Ryanotar 50 MG Injection

What is it prescribed for?

Intraabdominal Infection

Ryanotar 50 MG Injection is used in the treatment of Intra-abdominal infections like Peritonitis, Appendicitis, Intra-abdominal Abscess caused by E.coli, Streptococci, and Enterococci.

Skin and Structure Infection

Ryanotar 50 MG Injection is used in the treatment of Skin and Structure infection like cellulitis, wound infection and cutaneous abscess caused by Streptococcus pyogenes and Staphylococcus aureus.

Pneumonia

Ryanotar 50 MG Injection is used in the treatment of community acquired pneumonia which is a most common type of lung infection caused by Streptococcus pneumoniae and Haemophilus influenzae.

Food and Drug Administration (FDA) approved .Prescription Only (POM)

A Taj Pharma® India Product

TAJ
PHARMA®
A TAJ Enterprise

Do not store above 25°C. Keep container in the outer carton, in order to protect from light. Once reconstituted and diluted in the bag or other suitable infusion container (e.g. glass bottle), tigecycline should be used immediately.