www.tajpharma.com

SUSTRIXON® is the only underarm low testosterone (low T) treatment SUSTRIXON® is a prescription medication that contains testosterone and is used to treat adult men with low or no testosterone.

SUSTRIXON® is not intended for use in women or anyone under 18 years of age. It is not known if SUSTRIXON® is safe and effective in children younger than 18 years old. Improper use of SUSTRIXON® may affect bone growth in children.

NEW SUSTRIXON (testosterone) topical solution (Low Testosterone Treatment, Low T

(testosterone) topical solution (1)

Low Testosterone Treatment, Low T

SUSTRIXON * topical solution is an underarm testosterone treatment. Testosterone is a naturally occurring male hormone necessary for many processes in the body.

Indication: SUSTRIXON is used to treat adult males who have low or no testosterone. SUSTRIXON is not intended for use in women or anyone under 18 years of age. It is not known if SUSTRIXON is safe and effective in children younger than 18 years old. Improper use of SUSTRIXON may affect bone growth in children.

Important Safety Information for SUSTRIXON (testosterone) topical solution

What is the most important information I should know about SUSTRIXON?

SUSTRIXON can transfer from your body to others. This can happen if other people come into contact with the area where the SUSTRIXON was applied. Signs of puberty that are not expected (for example, pubic hair) have happened in young children who were accidentally exposed to testosterone through skin to skin contact with men using topical testosterone products like SUSTRIXON. Women and children should avoid contact with the unwashed or unclothed area where SUSTRIXON has been applied. If a woman or child makes contact with the application area, the contact area on the woman or child should be washed well with soap and water right away.

To lower the risk of transfer of SUSTRIXON from your body to others, follow these important instructions:

Apply SUSTRIXON only to your armpits.

Wash your hands right away with soap and water after applying SUSTRIXON.

After the solution has dried, cover the application area with clothing. Keep area covered until you have washed the application area well or have showered.

If you expect another person to have direct skin-to-skin contact with your armpits, first wash the application area well with soap and water.

Stop using SUSTRIXON and call your healthcare provider right away if you see any signs and symptoms in a child or a woman that may have occurred through accidental exposure to SUSTRIXON. Signs and symptoms in children may include enlarged penis or clitoris; early development of pubic hair; increased erections or sex drive; aggressive behavior. Signs and symptoms in women may include changes in body hair and a large increase in acne.

Who should not use SUSTRIXON?

Do not use SUSTRIXON if you:

have or might have prostate cancer

have breast cancer

are pregnant or may become pregnant or are breast-feeding. SUSTRIXON may harm your unborn or breast-feeding baby. Women who are pregnant or who may become pregnant should avoid contact with the area of skin where SUSTRIXON has been applied.

NEW SUSTRIXON (testosterone) topical solution (Low Testosterone Treatment, Low T