

# Tramadol Hydrochloride Injection 50mg/1ml, 100mg/2ml

## **1. Name of the medicinal product**

Tramadol Hydrochloride Injection  
50mg/1ml Taj Pharma

Tramadol Hydrochloride Injection  
100mg/2ml Taj Pharma

## **2. Qualitative and quantitative composition**

a) Tramadol Hydrochloride Injection  
50mg/1ml

Each ml contains:

Tramadol Hydrochloride	50mg
Water for injection	q.s

b) Tramadol Hydrochloride Injection  
100mg/2ml

Each ml contains:

Tramadol Hydrochloride	50mg
Water for injection	q.s

For the full list of excipients, see Section 6.1.

## **3. Pharmaceutical form**

Clear, colourless sterile solution for injection or infusion.

## **4. Clinical particulars**

### **4.1 Therapeutic indications**

Treatment of moderate to severe pain.

### **4.2 Posology and method of administration**

The dose should be adjusted to the intensity of the pain and the sensitivity of the individual patient. The lowest effective dose for analgesia should generally be selected.

The total daily dose of 400 mg tramadol

hydrochloride should not be exceeded, except in special clinical circumstances.

Unless otherwise prescribed, Tramadol hydrochloride 50 mg/ml solution for injection or infusion should be administered as follows:

### **Adults and adolescents above the age of 12 years:**

The usual dose is 50 or 100mg 4-6 hourly by the intravenous or intramuscular route. Dosage should be adjusted according to pain severity and response.

Intravenous injections must be given slowly over 2-3 minutes.

For post-operative pain administer an initial bolus of 100mg. During the 60 minutes following the initial bolus, further doses of 50mg may be given every 10-20 minutes, up to a total dose of 250mg including the initial bolus. Subsequent doses should be 50mg or 100mg 4-6 hourly up to a total daily dose of 400mg.

### **Children**

Tramadol hydrochloride 50 mg/ml solution for injection or infusion is not suitable for children below the age of 12 years.

### **Geriatric patients**

A dose adjustment is not usually necessary in elderly patients (up to 75 years) without clinically manifest hepatic or renal insufficiency. In elderly patients (over 75 years) elimination may be prolonged. Therefore, if necessary the dosage interval is to be extended according to the patient's requirements.

### **Renal Insufficiency/Dialysis and Hepatic Insufficiency**

In patients with renal and/or hepatic insufficiency the elimination of tramadol is delayed. In these patients prolongation of the dosage intervals should be carefully considered according to the patient's requirements.

#### **Method of administration**

Tramadol hydrochloride 50 mg/ml solution for injection or infusion may be administered intramuscularly, by slow intravenous injection, or diluted in solution (see Section 6.6) for administration by infusion or patient controlled analgesia.

#### **Duration of administration**

Tramadol hydrochloride 50 mg/ml solution for injection or infusion should under no circumstances be administered for longer than absolutely necessary. If long-term pain treatment with Tramadol hydrochloride 50 mg/ml solution for injection or infusion is necessary in view of the nature and severity of the illness, then careful regular monitoring should be carried out (if necessary with breaks in treatment) to establish whether and to what extent further treatment is necessary.

#### **4.3 Contraindications**

Tramadol hydrochloride 50 mg/ml solution for injection or infusion is contraindicated

- in patients who have previously shown hypersensitivity to the active substance tramadol or to any of the excipients listed in section 6.1.
- in patients suffering from acute intoxication with alcohol, hypnotics, analgesics, opioids, or psychotropic medicinal products.
- in patients who are receiving monoamine oxidase (MAO) inhibitors or who have taken

them within the last 14 days (see section 4.5)

- in patients with epilepsy not adequately controlled by treatment.
- for use in narcotic withdrawal treatment.

#### **4.4 Special warnings and precautions for use**

Tramadol hydrochloride 50 mg/ml solution for injection or infusion may only be used with particular caution in opioid-dependent patients, patients with head injury, shock, a reduced level of consciousness of uncertain origin, disorders of the respiratory centre or function, increased intracranial pressure.

In patients sensitive to opiates the product should only be used with caution.

Care should be taken when treating patients with respiratory depression, or if concomitant CNS depressant drugs are being administered (see section 4.5), or if the recommended dosage is significantly exceeded (see section 4.9) as the possibility of respiratory depression cannot be excluded in these situations.

Convulsions have been reported in patients receiving tramadol at the recommended dose levels. The risk may be increased when doses of tramadol exceed the recommended upper daily dose limit (400 mg). In addition, tramadol may increase the seizure risk in patients taking other medicinal products that lowers the seizure threshold (see section 4.5). Patients with epilepsy or those susceptible to seizures should only be treated with tramadol if there are compelling circumstances.

Tolerance, psychological and physical dependence may develop, especially after long-term use. In patients with a tendency to drug abuse or dependence, treatment with

tramadol should only be carried out for short periods under strict medical supervision.

Tramadol hydrochloride 50 mg/ml solution for injection or infusion is not a suitable substitute in opioid dependent patients. Although it is an opioid agonist, tramadol cannot suppress morphine withdrawal symptoms.

When a patient no longer requires therapy with tramadol, it may be advisable to taper the dose gradually to prevent symptoms of withdrawal.

#### CYP2D6 metabolism

Tramadol is metabolised by the liver enzyme CYP2D6. If a patient has a deficiency or is completely lacking this enzyme an adequate analgesic effect may not be obtained. Estimates indicate that up to 7% of the Caucasian population may have this deficiency. However, if the patient is an ultra-rapid metaboliser there is a risk of developing side effects of opioid toxicity even at commonly prescribed doses.

General symptoms of opioid toxicity include confusion, somnolence, shallow breathing, small pupils, nausea, vomiting, constipation and lack of appetite. In severe cases this may include symptoms of circulatory and respiratory depression, which may be life threatening and very rarely fatal. Estimates of prevalence of ultra-rapid metabolisers in different populations are summarised below:

Population	Prevalence %
African/Ethiopian	29%
African American	3.4% to 6.5%
Asian	1.2% to 2%
Caucasian	3.6% to 6.5%
Greek	6.0%
Hungarian	1.9%

Northern European	1% to 2%
-------------------	----------

#### Post-operative use in children

There have been reports in the published literature that tramadol given post-operatively in children after tonsillectomy and/or adenoidectomy for obstructive sleep apnoea, led to rare, but life threatening adverse events. Extreme caution should be exercised when tramadol is administered to children for post-operative pain relief and should be accompanied by close monitoring for symptoms of opioid toxicity including respiratory depression.

#### Children with compromised respiratory function

Tramadol is not recommended for use in children in whom respiratory function might be compromised including neuromuscular disorders, severe cardiac or respiratory conditions, upper respiratory or lung infections, multiple trauma or extensive surgical procedures. These factors may worsen symptoms of opioid toxicity.

#### Risk from concomitant use of sedative medicines such as benzodiazepines or related drugs

Concomitant use of Tramadol hydrochloride 50 mg/ml solution for injection or infusion and sedative medicines such as benzodiazepines or related drugs may result in sedation, respiratory depression, coma and death. Because of these risks, concomitant prescribing with these sedative medicines should be reserved for patients for whom alternative treatment options are not possible. If a decision is made to prescribe Tramadol hydrochloride 50 mg/ml solution for injection or infusion concomitantly with sedative medicines, the lowest effective dose should be used, and the duration of treatment should be as short as possible.

The patients should be followed closely for signs and symptoms of respiratory depression and sedation. In this respect, it is strongly recommended to inform patients and their caregivers to be aware of these symptoms (see section 4.5).

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Tramadol hydrochloride 50 mg/ml solution for injection or infusion should not be combined with MAO inhibitors (see section 4.3).

In patients treated with MAO inhibitors in the 14 days prior to the use of the opioid pethidine, life-threatening interactions on the central nervous system, respiratory and cardiovascular function have been observed. The same interactions with MAO inhibitors cannot be ruled out during treatment with Tramadol hydrochloride 50 mg/ml solution for injection or infusion.

The results of pharmacokinetic studies have so far shown that on the concomitant or previous administration of cimetidine (enzyme inhibitor) clinically relevant interactions are unlikely to occur. Simultaneous or previous administration of carbamazepine (enzyme inducer) may reduce the analgesic effect and shorten the duration of action.

Tramadol can induce convulsions and increase the potential for selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants, antipsychotics and other seizure threshold-lowering medicinal products (such as bupropion, mirtazapine, tetrahydrocannabinol) to cause convulsions.

Concomitant therapeutic use of tramadol and serotonergic drugs, such as selective serotonin reuptake inhibitors (SSRIs),

serotonin-norepinephrine reuptake inhibitors (SNRIs), MAO inhibitors (see section 4.3), tricyclic antidepressants and mirtazapine may cause serotonin toxicity. Serotonin syndrome is likely when one of the following is observed:

- Spontaneous clonus
- Inducible or ocular clonus with agitation or diaphoresis
- Tremor and hyperreflexia
- Hypertonia and body temperature > 38 °C and inducible or ocular clonus.

Withdrawal of the serotonergic drugs usually brings about a rapid improvement. Treatment depends on the type and severity of the symptoms.

Caution should be exercised during concomitant treatment with tramadol and coumarin derivatives (e.g. warfarin) due to reports of increased INR with major bleeding and ecchymoses in some patients.

Other active substances known to inhibit CYP3A4, such as ketoconazole and erythromycin, might inhibit the metabolism of tramadol (N-demethylation) probably also the metabolism of the active O-demethylated metabolite. The clinical importance of such an interaction has not been studied (see section 4.8).

In a limited number of studies, the pre- or postoperative application of the antiemetic 5-HT<sub>3</sub> antagonist ondansetron increased the requirement of tramadol in patients with postoperative pain.

The administration of Tramadol hydrochloride 50 mg/ml solution for injection or infusion with other centrally depressant medicinal products, including

alcohol, may potentiate the CNS effects (see section 4.8).

#### Sedative medicines such as benzodiazepines or related drugs

The concomitant use of opioids with sedative medicines such as benzodiazepines or related drugs increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect. The dose and duration of concomitant use should be limited (see section 4.4).

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

Animal studies with tramadol revealed at very high doses effects on organ development, ossification and neonatal mortality. Tramadol crosses the placenta. There is inadequate evidence available on the safety of tramadol in human pregnancy. Therefore Tramadol hydrochloride 50 mg/ml solution for injection or infusion should not be used in pregnant women.

Tramadol - administered before or during birth - does not affect uterine contractility. In neonates it may induce changes in the respiratory rate which are usually not clinically relevant. Chronic use during pregnancy may lead to neonatal withdrawal symptoms.

#### Breast-feeding

Approximately 0.1 % of the maternal dose is excreted into the milk. In the immediate post-partum period, for maternal oral daily dosage up to 400 mg, this corresponds to a mean amount of tramadol ingested by breast-fed infants of 3% of the maternal weight-adjusted dosage. For this reason tramadol should not be used during lactation or alternatively, breast-feeding should be discontinued during treatment with

tramadol. Discontinuation of breast-feeding is generally not necessary following a single dose of tramadol.

#### Fertility

Post marketing surveillance does not suggest an effect of tramadol on fertility. Animal studies did not show an effect of tramadol on fertility.

### **4.7 Effects on ability to drive and use machines**

Even when taken according to instructions, Tramadol hydrochloride 50 mg/ml solution for injection or infusion may cause effects such as somnolence and dizziness and therefore may impair a patient's ability to drive safely or operate machinery. This applies particularly in conjunction with alcohol and other psychotropic substances. Patients should, therefore, not drive or operate machinery.

*This class of medicine is in the list of drugs included in regulations under 5a of the Road Traffic Act 1988. When prescribing this medicine, patients should be told:*

- *The medicine is likely to affect your ability to drive*
- *Do not drive until you know how the medicine affects you*
- *It is an offence to drive while under the influence of this medicine*
- *However, you would not be committing an offence (called 'statutory defence') if:*
  - o *The medicine has been prescribed to treat a medical or dental problem and*
  - o *You have taken it according to the instructions given by the prescriber and in the information provided with the medicine and*

o It was not affecting your ability to drive safely

#### **4.8 Undesirable effects**

Rapid intravenous administration may be associated with a higher incidence of adverse effects and therefore should be avoided.

The most commonly reported adverse drug reactions are nausea and dizziness, both occurring in more than 10% of patients.

The frequencies are defined as follows:

*Very common:*  $\geq 1/10$

*Common:*  $\geq 1/100$ ,  $< 1/10$

*Uncommon:*  $\geq 1/1000$ ,  $< 1/100$

*Rare:*  $\geq 1/10\ 000$ ,  $< 1/1000$

*Very rare:*  $< 1/10\ 000$

*Not known:* cannot be estimated from the available data

#### Cardiovascular disorders:

*uncommon:* cardiovascular regulation (palpitation, tachycardia). These adverse reactions may occur especially on intravenous administration and in patients who are physically stressed.

*rare :* bradycardia

#### Investigations:

Rare: increase in blood pressure

#### Vascular disorders:

*Uncommon:* cardiovascular regulation (postural hypotension or cardiovascular collapse). These adverse reactions may occur especially on intravenous administration and in patients who are physically stressed.

#### Metabolism and nutrition disorders:

*Rare:* changes in appetite

*Not known:* hypoglycaemia

#### Respiratory, thoracic and mediastinal disorders:

*Rare:* respiratory depression, dyspnoea

If the recommended doses are considerably exceeded and other centrally depressant substances are administered concomitantly (see section 4.5), respiratory depression may occur.

Worsening of asthma has been reported, though a causal relationship has not been established.

#### Nervous system disorders:

*very common :* dizziness

*common :* headache, somnolence

*rare :* changes in appetite, paraesthesia, tremor, respiratory depression, epileptiform convulsions, involuntary muscle contractions, abnormal coordination, syncope.

*not known:* speech disorders

Convulsions occurred mainly after administration of high doses of tramadol or after concomitant treatment with medicinal products which can lower the seizure threshold (see sections 4.4 and 4.5).

#### Psychiatric disorders:

*rare :* hallucinations, confusion, sleep disturbance, delirium, anxiety and nightmares. Psychological adverse reactions may occur following administration of Tramadol hydrochloride 50 mg/ml solution for injection or infusion which vary individually in intensity and nature

(depending on personality and duration of treatment). These include changes in mood (usually elation, occasionally dysphoria), changes in activity (usually suppression, occasionally increase) and changes in cognitive and sensorial capacity (e.g. decision behaviour, perception disorders). Dependence may occur.

Symptoms of withdrawal reactions, similar to those occurring during opiate withdrawal, may occur as follows: agitation, anxiety, nervousness, insomnia, hyperkinesia, tremor and gastrointestinal symptoms. Other symptoms that have very rarely been seen with tramadol discontinuation include: panic attacks, severe anxiety, hallucinations, paraesthesias, tinnitus and unusual CNS symptoms (i.e. confusion, delusions, depersonalisation, derealisation, paranoia).

Eye disorders:

*rare* : miosis, mydriasis, blurred vision

Gastrointestinal disorders:

*very common* : nausea

*common* : vomiting, constipation, dry mouth

*uncommon* : retching; gastrointestinal irritation (a feeling of pressure in the stomach, bloating), diarrhoea

Skin and subcutaneous tissue disorders:

*common* : sweating

*uncommon* :dermal reactions (e.g. pruritus, rash, urticaria)

Musculoskeletal and connective tissue disorders:

*rare* : motorial weakness

Hepatobiliary disorders:

In a few isolated cases an increase in liver enzyme values has been reported in a temporal connection with the therapeutic use of tramadol.

Renal and urinary disorders:

*rare* : micturition disorders (difficulty in passing urine, dysuria and urinary retention)

Immune system disorders:

*Rare*: allergic reactions (e.g. dyspnoea, bronchospasm, wheezing, angioneurotic oedema) and anaphylaxis

General disorders:

*common* : fatigue

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important.

## 4.9 Overdose

Symptoms

In principle, on intoxication with tramadol symptoms similar to those of other centrally acting analgesics (opioids) are to be expected. These include in particular miosis, vomiting, cardiovascular collapse, consciousness disorders up to coma, convulsions and respiratory depression up to respiratory arrest.

Treatment

The general emergency measures apply. Keep open the respiratory tract (aspiration!), maintain respiration and circulation depending on the symptoms. The antidote for respiratory depression is naloxone. In animal experiments naloxone had no effect on convulsions. In such cases diazepam should be given intravenously.

In case of intoxication orally, gastrointestinal decontamination with activated charcoal or by gastric lavage is only recommended within 2 hours after tramadol intake. Gastrointestinal decontamination at a later time point may be useful in case of intoxication with exceptionally large quantities.

Tramadol is minimally eliminated from the serum by haemodialysis or haemo-filtration. Therefore treatment of acute intoxication with Tramadol hydrochloride 50 mg/ml solution for injection or infusion with haemodialysis or haemofiltration alone is not suitable for detoxification.

## **5. Pharmacological properties**

### **5.1 Pharmacodynamic properties**

Tramadol is a centrally acting analgesic which possesses opioid agonist properties. Tramadol consists of two enantiomers, the (+)-isomer is predominantly active as an opioid with preferential activity for the  $\mu$ -receptor. The (-)-isomer potentiates the analgesic effect of the (+)-isomer and is active as an inhibitor of noradrenaline and serotonin uptake thereby modifying the transmission of pain impulses.

Tramadol also has an antitussive action. At the recommended dosages, the effects of tramadol given orally on the respiratory and cardiovascular systems appear to be clinically insignificant. The potency of tramadol is reported to be  $1/10^{\text{th}}$  to  $1/6^{\text{th}}$  that of morphine.

### **Paediatric population**

Effects of enteral and parenteral administration of tramadol have been investigated in clinical trials involving more than 2000 paediatric patients ranging in age from neonate to 17 years of age. The indications for pain treatment studied in

those trials included pain after surgery (mainly abdominal), after surgical tooth extractions, due to fractures, burns and traumas as well as other painful conditions likely to require analgesic treatment for at least 7 days.

At single doses of up to 2 mg/kg or multiple doses of up to 8 mg/kg per day (to a maximum of 400 mg per day) efficacy of tramadol was found to be superior to placebo, and superior or equal to paracetamol, nalbuphine, pethidine or low dose morphine. The conducted trials confirmed the efficacy of tramadol. The safety profile of tramadol was similar in adult and paediatric patients older than 1 year (see section 4.2).

### **5.2 Pharmacokinetic properties**

#### **a) General**

The mean absolute bioavailability after intramuscular administration was found to be 100%.

The distribution of tramadol following intravenous administration is rapid and in two phases with different half-lives of  $0.31 \pm 0.17$  hours (initial rapid phase) and  $1.7 \pm 0.4$  hours (slower phase) respectively.

After intravenous administration of 100 mg tramadol, the serum concentration was  $613 \pm 221$  ng/ml at 15 minutes post dosing and  $409 \pm 79$  ng/ml at 2 hours post dosing. Tramadol has a high tissue affinity with an apparent volume of distribution of 203 L after intravenous dosing in healthy volunteers.

Tramadol undergoes hepatic metabolism with approximately 85% of an intravenous dose being metabolised in young healthy volunteers. In humans tramadol is mainly metabolised by means of N- and O-demethylation and conjugation of the O-



demethylation products with glucuronic acid. Only O-desmethyltramadol is pharmacologically active. There are considerable interindividual quantitative differences between the other metabolites. So far, eleven metabolites have been found in the urine. Animal experiments have shown that O-desmethyltramadol is more potent than the parent substance by the factor 2-4. Its half life  $t_{1/2\beta}$  (6 healthy volunteers) is 7.9 h (range 5.4-9.6 h) and is approximately that of tramadol.

The inhibition of one or both cytochrome P450 isoenzymes, CYP3A4 and CYP2D6 involved in the metabolism of tramadol, may affect the plasma concentration of tramadol or its active metabolite.

Tramadol is essentially excreted via the kidneys. The mean elimination half-life of tramadol following intravenous administration is 5-6 hours. Total clearance of tramadol was 28.0 L/h following intravenous administration.

#### **b) Characteristics in patients**

Effect of age: Tramadol pharmacokinetics show little age-dependence in volunteers up to the age of 75 years. In volunteers aged over 75 years, the terminal elimination half-life was  $7.0 \pm 1.6$  h compared to  $6.0 \pm 1.5$  h in young volunteers after oral administration. Effect of hepatic or renal impairment: As both tramadol and its pharmacologically active metabolite, O-desmethyl tramadol, are eliminated both metabolically and renally, the terminal half-life of elimination ( $t_{1/2}$ ) may be prolonged in patients with hepatic or renal dysfunction. However, the increase in  $t_{1/2}$  is relatively small if either excretory organ is functioning normally. In liver cirrhosis patients, the mean  $t_{1/2}$  of tramadol was  $13.3 \pm 4.9$  hours. In patients with renal failure (creatinine

clearance  $< 5$  mL/min) the  $t_{1/2}$  of tramadol was  $11.0 \pm 3.2$  hours and that of M1 was  $16.9 \pm 3.0$  hours. Extreme values observed to date are 22.3 hours (tramadol) and 36.0 hours (M1) in liver cirrhosis patients and 19.5 hours (tramadol) and 43.2 hours (M1) in renal failure patients.

#### Paediatric population

The pharmacokinetics of tramadol and O-desmethyltramadol after single-dose and multiple-dose oral administration to subjects aged 1 year to 16 years were found to be generally similar to those in adults when adjusting for dose by body weight, but with a higher between-subject variability in children aged 8 years and below.

In children below 1 year of age, the pharmacokinetics of tramadol and O-desmethyltramadol have been investigated, but have not been fully characterized. Information from studies including this age group indicates that the formation rate of O-desmethyltramadol via CYP2D6 increases continuously in neonates, and adult levels of CYP2D6 activity are assumed to be reached at about 1 year of age. In addition, immature glucuronidation systems and immature renal function may result in slow elimination and accumulation of O-desmethyltramadol in children under 1 year of age.

#### **5.3 Preclinical safety data**

On repeated oral and parenteral administration of tramadol for 6 - 26 weeks in rats and dogs and oral administration for 12 months in dogs, haematological, clinico-chemical and histological investigations showed no evidence of any substance-related changes. Central nervous manifestations only occurred after high doses considerably above the therapeutic range: restlessness, salivation, convulsions, and reduced weight gain. Rats and dogs

tolerated oral doses of 20 mg/kg and 10 mg/kg body weight respectively, and dogs rectal doses of 20 mg/kg body weight without any reactions.

In rats tramadol dosages from 50 mg/kg/day upwards caused toxic effects in dams and raised neonate mortality. In the offspring retardation occurred in the form of ossification disorders and delayed vaginal and eye opening. Male fertility was not affected. After higher doses (from 50 mg/kg/day upwards) females exhibited a reduced pregnancy rate. In rabbits there were toxic effects in dams from 125 mg/kg upwards and skeletal anomalies in the offspring.

In some in-vitro test systems there was evidence of mutagenic effects. In-vivo studies showed no such effects. According to knowledge gained so far, tramadol can be classified as non-mutagenic.

Studies on the tumorigenic potential of tramadol hydrochloride have been carried out in rats and mice. The study in rats showed no evidence of any substance-related increase in the incidence of tumours. In the study in mice there was an increased incidence of liver cell adenomas in male animals (a dose-dependent, non-significant increase from 15 mg/kg upwards) and an increase in pulmonary tumours in females of all dosage groups (significant, but not dose-dependent).

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

Sodium acetate trihydrate

Water for injections

### **6.2 Incompatibilities**

Precipitation will occur if Tramadol hydrochloride 50 mg/ml solution for

injection or infusion is mixed in the same syringe with injections of diazepam, diclofenac sodium, indomethacin, midazolam and piroxicam.

### **6.3 Shelf life**

36 months. From a microbiological point of view, the product should be used immediately. If not used-immediately, in-use storage times and conditions prior to use are the responsibility of the user.

### **6.4 Special precautions for storage**

Do not store above 30°C.

### **6.5 Nature and contents of container**

Colourless glass ampoule containing either 1 ml or 2 ml of injection solution. Ampoules are contained in a pre-fabricated blister strip, which is enclosed in a cardboard outer carton. Cartons contain either 1, 5 or 10 ampoules.

### **6.6 Special precautions for disposal and other handling**

Based on described observations Tramadol 50 mg/ml solution for injection or infusion is physically and chemically compatible for up to 24 hours with water for injections (WFI), sodium chloride solution for injection 0.9% and glucose solution for injection 5% at given concentrations. Obtained results show no necessity for cold storage.

The prepared infusion solution should be made immediately before use.

## **7. Manufactured in India by:**

**TAJ PHARMACEUTICALS LTD.**

Mumbai, India

Unit No. 214.Old Bake House,

Maharashtra chambers of Commerce Lane,

Fort, Mumbai - 400001

at:Gujarat, INDIA.

Customer Service and Product Inquiries:



1-800-TRY-FIRST (1-800-222-434 & 1-800-222-825)

Monday through Saturday 9:00 a.m. to 7:00 p.m. EST

E-mail: [tajgroup@tajpharma.com](mailto:tajgroup@tajpharma.com)